

Magnetic Field Diffusion and the Formation of Circumstellar Disks

Shantanu Basu¹, Wolf B. Dapp^{1,2}, and Matthew W. Kunz³

¹Department of Physics and Astronomy, University of Western Ontario, London, Ontario, N6A 3K7, Canada

²Jülich Supercomputing Centre, Institute for Advanced Simulation, FZ Jülich, Germany

³Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA; Einstein Postdoctoral Fellow

E-mail: basu@uwo.ca

Abstract. A non-ideal MHD collapse calculation employing the axisymmetric thin-disk approximation is used to resolve cloud core collapse down to the scales of the second (stellar) core. Rotation and a magnetic braking torque are included in the model, and the partial ionization resulting in ambipolar diffusion and Ohmic dissipation is calculated from a detailed chemical network. We find that a centrifugal disk can indeed form in the earliest stage of star formation, due to a shut-off of magnetic braking caused by magnetic field diffusion in the first core region. Thus, there is no catastrophic magnetic braking in a model with realistic non-ideal MHD.

1. Results and Model Description

Fig. 1 shows the magnetic field lines within the inner 10 AU at the end of the simulation in two different cases. A second (stellar) core has just formed in both cases, after second collapse of the first hydrostatic core. The second core is centered at the origin ($r = 0, z = 0$) and the thin-disk model extends in the equatorial plane. The magnetic field lines are obtained above and below the midplane using the force-free and current-free approximation, using the currents in the midplane as a source term.

The flux-frozen model (dashed lines) shows extreme pinching of the magnetic field lines, into nearly a split-monopole configuration. This is due to the dragging-in of field lines in the flux-freezing limit. The extreme flaring of magnetic field lines results in catastrophic magnetic braking in this model. Magnetic field lines that tie small radii near the protostar with a much larger lever arm far above the disk can very efficiently extract angular momentum from the midplane region. No centrifugally-supported disk is able to form in this case, and radial infall continues onto the protostar. Conversely, the non-ideal MHD model (solid lines) shows that the same field lines straighten out significantly on the small scales shown in the figure (the full model extends to about 10^4 AU in radius). The straightening of field lines on small scales shuts down the efficiency of magnetic braking, and a disk of radius $\approx 10 R_{\odot}$ is formed when the protostar




Figure 1. Magnetic field lines. The box has dimensions of 10 AU on each side, and the newly formed second core is located at the origin. The dashed lines represent a flux-frozen model, while the solid lines show the same field lines for a model including non-ideal MHD effects. Note the extreme flaring of the magnetic field in the flux-frozen model (leading to catastrophic magnetic braking) and the straightening out of field lines in the non-ideal MHD model.

has mass $\approx 10^{-3} M_{\odot}$ and has just begun the accretion process. Based on the angular momentum content of the core at this time, we estimate that direct infall will lead to a small disk (radius < 10 AU) for $\approx 4 \times 10^4$ yr, representing the early Class 0 phase.

This result extends previous work on disk formation that employed a simplified parametrization of just Ohmic dissipation [1]. Here, we use a full chemical network model to obtain partial ionization at each location in the cloud and thereby calculate local coefficients of both ambipolar diffusion and Ohmic dissipation. We adopt the model of [2] in this regard. A detailed presentation of our results can be found in [3].

References

- [1] Dapp, W. B., & Basu, S. 2010, *A & A*, 521, L56
- [2] Kunz, M. W., & Mouschovias, T. Ch. 2009, *Astrophys. J.*, 693, 1895
- [3] Dapp, W. B., Basu, S., & Kunz, M. W. 2012, arXiv:1112.3801